

EVALUATION OF ANTICANCER ACTIVITY OF CHLORELLA VULGARIS AGAINST HUMAN BREAST ADENOCARCINOMA CELL LINE (MCF7)

Prakash B¹, Virgin Nisha M¹, Miller Samson S¹, Kavitha R¹, Ashokkumar L² and Jegadeeshkumar D³

¹Department of Biotechnology, Vivekanandha College of Arts and Sciences for Women (Autonomous), Elayampalyam, Tiruchengode -637205, Namakkal, Tamilnadu, India.

²Department of Microbiology, Vivekanandha College of Arts and Sciences for Women (Autonomous), Elayampalyam, Tiruchengode -637205, Namakkal, Tamilnadu, India.

³Chromopark Research Centre, Namakkal, Tamilnadu, India

Abstract

Cancer is an increasing trend and successful therapies are inadequate to approach these malignancies. Normally rapidly dividing cells are controlled by anti-cancer drugs, but the normal cells are also affected and pattern in which is determined the side effects. A great number of antitumor compounds are natural products or their derivatives, mainly produced by blue-green algae. The methanol extract of *Chlorella vulgaris* was reported to possess anticancer effects against human tumor cell lines. The results showed the inhibition of in vitro proliferation of human tumor cell lines against MCF7 cell lines. Totally seven types of Concentration of plant extracts were subjected to anticancer activity, among them 7 concentrations were exhibited anticancer activity, the fifty percentage of cell death occur when using 31.2µg of plant extract. In our present investigation shows that the phytochemical's of *Chlorella vulgaris*, may be responsible for displaying anticancer activities..

Keywords *Chlorella vulgaris*, MCF7, Cancer.

INTRODUCTION

Malignant tumor incidences are in increasing trend and effective therapies are inadequate to approach these malignancies. Normally rapidly dividing cells are controlled by anti-cancer drugs, but the normal cells are also affected and pattern in which is determined the side effects. The behavior in which the disparate cells are concerned determines the side effects of the all by one lonesome drug. These side effects may be minimized by improving and new remedial preparations. These drugs could be of ethno botanical origin. Auspiciously numerous preceding readings have shown that the anticancer activities of non-toxic biological macromolecules are higher than conventional chemotherapy drugs. Marine algae is obliged as significant sources of natural bioactive substances and there has now emerged a new proclivity towards isolating and identifying such compounds and constituents from algae. This review article has poised studies about algal anticancer agents [1]. In 2016, Mohd *et al*, determine the anticancer activity of *Chlorella vulgaris* against breast

cancer cell lines. The tests were carried out by means of MTT assay (a colorimetric assay) to determine cell viability by assessing the values of IC₅₀. The result showed that the MCF7 was the cancer cell lines that reach 50% of inhibition by algae.

The extraction by chloroform of *Chlorella vulgaris* indicates that there was 50% inhibition (IC₅₀) of cells on MCF7 much higher compare to the ethanol extraction of both spirulina and chlorella with at 89 µg/ml of extract. In recently [2] also observed the anticancer activity against human breast cancer cell lines. They were observed the anticancer activity by MTT assay and 50% of cell death with 28 µg/ml of extract. Therefore, the present study is to investigate *Chlorella vulgaris* (*C. vulgaris*), a kind of freshwater green microalgae, which evaluates the anticancer activity against MCF7.

MATERIALS AND METHODS

The sample microalgae species *Chlorella vulgaris* sp. was obtained from Royal Research Centre, Chennai. The algae was cultured on using Bold Basal Medium (BBM). The cultures were grown at 24±1°C in a thermostatically controlled room with cool white fluorescent lamps at 2-3 weeks. After incubation algal growth was measured by using UV-VIS spectrophotometer at 680nm. Ten ml from cultures were filtered under vacuum using filter membrane (0.45µm) and washed several times with distilled

Address Correspondence @ Department of Biotechnology, Vivekanandha College of Arts and Sciences for Women (Autonomous), Elayampalyam-637205, Tiruchengode, Namakkal, Tamilnadu, India. Cell: + 91 94432 04101 Email: prakazbt@gmail.com



water. Then, the algae cells were dried at 80°C for 30min and weighed.

Preparation of algae extracts

Extraction was performed using the solvents of different polarity: acetone and Methanol. Five grams of the algal dry powder were suspended in the solvents at the ratio of 6:1 (v/w), left at room temperature for 48 h and then, were homogenized for 30 min. Finally, the suspensions were centrifuged at 3000 g for 10 min and the supernatants were filtered and concentrated. The extracts were kept in dark at 4°C until use. Followed by the preliminary phytochemical screening was performed [3].

Cytotoxicity assay

Antitumor activity assay of the extracts was determined by Human Breast Cancer (MCF7). The cell line was obtained from the NCCS, Pune. The cells were grown in RPMI- 1640 medium supplemented with 10% inactivated fetal calf serum and 50µg/ml gentamycin. The cells were maintained at 37°C in a humidified with 5% CO₂ and were subcultures two to three times a week, according to Mohd Syahril[3].

Trypan blue exclusion test

To detect the cell viability trypan blue exclusion test was chosen. The cancer cells were seeded at a density of 6×10⁵ cells/well with different concentration of extract at 37°C in the presence of 5% CO₂. After 72 hour, 20 µl of medium and equal volume of trypan blue were mixed and viable and dead cells were counted by haemocytometer.

MTT Assay

MTT assay as a proper cytotoxicity test was used in this study. Briefly, 10 µl of MTT stock solution [5 mg/ml in phosphate buffer saline] was added to 90 µl medium of wells. The microplate was incubated at 37°C for 4 hours and then, the optical density of each well was read by the ELISA reader (TECAN, Switzerland) in 540nm


Calculation

$$\% \text{ cell viability} = \frac{A_{540} \text{ of treated cells}}{A_{540} \text{ of control cells}} \times 100\%$$

RESULTS AND DISCUSSION

In this study, the crude methanol extract revealed the presence of alkaloids, Carbohydrate, Saponins, Terpinoids and Quinines but Flavonoids Phenols, Steroids Tannins and proteins show negative results. The methanol extract of *Chlorella vulgaris* was reported to possess anticancer effects against human tumor cell lines. The results showed the inhibition of in vitro proliferation of human tumor cell lines against MCF7 cell lines. This parameter was carryout with Methyl thiazolyl tetrazolium (MTT) assay. Totally seven types of Concentration of plant extracts were

subjected to anticancer activity, among them 7 concentrations exhibited anticancer activity, the fifty percentage of cell death occur when using 31.2µg of plant extract (Graph.1).



Graph 1: Percentage of cell viability of MCF7 cells

Chemotherapy is one of the main treatments used to cure cancer. Besides that, a group of drugs are used to kill or inhibit the growth of cancer cells. These drugs are associated with toxicity and side effects of chemotherapeutic drugs include hair loss, mouth sores, diarrhea, nausea and vomiting, loss of appetite and fatigue.




Figure.1: Microscopic examination of anticancer activity

Hence new anticancer agents should be investigated from various resources. A great number of antitumor compounds are natural products or their derivatives, mainly produced by blue-green algae [2]. *Chlorella vulgaris* was shown to have chemo preventive effect in induced liver cancer and breast cancer rats and possess a glycoprotein with antitumor effects [4,5]. In this current study, the methanol extract of *Chlorella vulgaris* was subjected to the anticancer activity against the MCF7 cancer cell line. The fifty percentage of cell death occurs when using 31.2µg of plant extract (Figure 1). The previous report emphasized the importance of strain selection as activity varied between strains of the same species [6,7]. Similar results were observed from previous study of Jayashree *et al.*, (2016) [8]. They were also observed the 50% of cell death from 28.1 µg of Methanolic *Chlorella vulgaris* extract against MCF 7.

CONCLUSION

Phytochemicals of the algae may responsible for displaying antimicrobial and anticancer activities. It can be pointed out from this study that microalgae have the potential to be further developed as an anticancer agent taking into account that it is a product of nature. However, further investigation is required for a complete analysis of its anticancer properties such as resolve the mechanisms of cell death caused by the algal extracts.

REFERENCES

- [1] **Nadia Sharif, NeelmaMunir, FaizaSaleem, FarheenAslam and ShaguftaNaz** (2014) "Prolific Anticancer Bioactivity of Algal Extracts (Review)". American Journal of Drug Delivery and Therapeutics. 1(2):060-072.
- [2] **Paterson MC, Dietrich KD, Danyluk J, Paterson AHG, Lees AW, Jamil N, Hanson J, Jenkins H, Krause BE, McBlain WA, Slamon DJ and Fournery RM** (1991) "Correlation between c-erbB-2 Amplification and Risk of Recurrent Disease in Node-negative Breast Cancer" Cancer Res, 51(2):556-567
- [3] **Mohd Syahril MZ, Roshani O, Nur hasyimah R, Mohamad Hafiz MS, Sharida MD, Ahmed HY** (2012) Screening of anticancer activities of crude extracts of unicellular green algae (*Chlorella vulgaris*) and filamentous blue green algae (*Spirulina platensis*) on selected cancer cell lines. Journal of Academia Univeriti Teknologi MARA Negeri Sembilan, 2:38-42.
- [4] **Solomon Charles Ugochukwu, Arukwe Uche I and Onuohalfeanyi** (2013) Preliminary phytochemical screening of different solvent extracts of stem bark and roots of *Denetiatripetala*, Asian Journal of Plant Science and Research, 3(3):10-13.
- [5] **Sulaiman S, Shamaan NA, Ngah WZW and Yusof YAM** (2006) Chemopreventive effect of *Chlorella vulgaris* in choline deficient diet and ethionine induced liver carcinogenesis in rats, Int. J. Cancer Res., 2: 234-241.
- [6] **Ruhul Amin ARM, Omer Kucuk, Fadlo RKhuri, Dong M Shin,** (2009) Perspectives for Cancer Prevention With Natural Compounds, Clinical Oncology 27(16): 2712-2725.
- [7] **Ordog V, Stirk WA, Lenobel R, Bancirova M, Strand M and Van Standen J,** (2004) Screening microalgae for some potentially useful agricultural and pharmaceutical secondary metabolites. J. Applied Phycol, 16:309-314.
- [8] **Jayashri Apurva, Dipali Barate, Musaddiq**(2013) Biofilm Forming Abilities and Antibiotic Susceptibility Pattern of Clinical Isolates of *Staphylococcus aureus*. Indian journal of applied research, 3(5):41-44.

Citation: Prakash B, Virgin Nisha M, Miller Samson S, Kavitha R, Ashokkumar L and Jegadeeshkumar D (2017) Evaluation of anticancer activity of *Chlorella vulgaris* against human breast adenocarcinoma cell line (MCF7), Int J Adv Interdis Res, 4 (4): 1-3.

Received: Oct 17, 2017 | **Revised:** Nov 15, 2017 | **Accepted:** Nov 16, 2017 | **Published:** Nov 26, 2017

License: Prakash et al (2017). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

